

Quark number scaling of v_2 in transverse kinetic energy and its implications for coalescence models

Jiangyong Jia^{1,3} and Chun Zhang²

¹*Columbia University, New York, NY 10027 and Nevis Laboratories, Irvington, NY 10533, USA*

²*Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*

³*Department of Chemistry, Stony Brook University, Stony Brook, NY 11794, USA*

(Dated: February 2, 2008)

We find that a simple extension of the coalescence model is sufficient to incorporate the perfect quark number scaling behavior of the elliptic flow in transverse kinetic energy, recently discovered by the PHENIX Collaboration. The flavor dependence of the elliptic flow can be consistently described in the low and intermediate p_T if the transverse kinetic energy is conserved in the $2 \rightarrow 1$ or $3 \rightarrow 1$ parton coalescence process at the hadronization. Thus suggesting the quark coalescence as a possible hadronization mechanism at low p_T as well.

PACS numbers: 25.75.Dw

Quark recombination or coalesce models [1, 2, 3, 4, 5] have been proposed to explain the large baryon to meson ratio [6, 7] and the number of constituent quark scaling (NCQ-scaling) of the hadron v_2 in p_T [7, 8] seen in RHIC data. Since coalescence models explicitly assume partonic degree of freedom, their apparent, albeit qualitative, success in describing the data at intermediate p_T is a strong support of the idea that the partonic matter has been created in Au-Au collisions at RHIC.

Coalescence models calculate the meson production via the convolution of Wigner functions for constituent quarks and the meson [2, 3, 4, 5]:

$$\frac{d^3 N}{dp^3} = \int \prod_{i=1}^2 d^3 x_i d^3 p_i F_M(x_1, p_1, x_2, p_2) W_M(p, p_1, p_2, x_1, x_2) \quad (1)$$

where F is the joint q and $\bar{q}$ phase space distribution, W_M is the meson Wigner function, which is typically approximated by

$$W_M = \Phi_M(x_1 - x_2, p_1 - p_2) \delta(p_T - p_{T,1} - p_{T,2}) \quad (2)$$

The δ function enforce the momentum conservation. It is straightforward to generalize these formula for baryons.

There are several implicit assumptions in coalescence models. Only valence quark degrees of freedom are assumed in the coalescence calculation, and the dynamic gluon contribution (appears as higher Fock state) [9] could lead to a 20% correction to the NCQ-scaling. There is a unitarity problem at low p_T [2, 5] due the quadratical dependence of the meson yield on the number of quarks, although this can be reconciled when one take into a finite freeze out time and no quarks are used in more than one hadron [10]. In addition, coalescence process reduce the entropy. This can be partially resolved by including resonance decay of unstable hadrons [11].

As Eq.1 indicates, the level of NCQ-scaling depends on both the final state parton phase space distribution, described by F_M , and the hadronization process which is encoded in W_M . The constituent quark phase space

distributions are driven by the space-time evolution of the partonic matter, which is typically described by hydrodynamics or parton cascade. Coalescence models explicitly assume that F_M can be factorized into the single quark distribution function, $F_M = f(x_1, p_1)f(x_2, p_2)$, with identical anisotropic phase space distribution (only v_2 is considered here.)

$$f(x, p) = n(x, p) (1 + 2v_{2,q}(x, p)\cos(2\phi)). \quad (3)$$

For a naive coalescence model where q and $\bar{q}$ are assumed to be co-moving : $\Phi_M \propto \delta^3(x_1 - x_2)\delta^3(p_1 - p_2)$, the spatially averaged v_2 for quark and meson is [12]:

$$\overline{v_{2,q}} = \frac{\int d^3 x d\phi f(x, \frac{p}{2}) \cos(2\phi)}{\int d^3 x d\phi f(x, \frac{p}{2})} \quad (4)$$

$$= \frac{\langle n(x, \frac{p}{2}) v_{2,q}(x, \frac{p}{2}) \rangle_x}{\langle n(x, \frac{p}{2}) \rangle_x}$$

$$\overline{v_{2,M}} = \frac{\int d^3 x d\phi f^2(x, \frac{p}{2}) \cos(2\phi)}{\int d^3 x d\phi f^2(x, \frac{p}{2})} \quad (5)$$

$$= \frac{2 \langle n^2(x, \frac{p}{2}) v_{2,q}(x, \frac{p}{2}) \rangle_x}{\langle n^2(x, \frac{p}{2}) (1 + 2v_{2,q}^2(x, \frac{p}{2})) \rangle_x}$$

where $\langle A(x) \rangle_x = \int d^3 x A(x)$. Because quark density $n(x, p)$ and parton anisotropy $v_{2,q}(x, p)$ could depend on quark local coordinate and momentum, NCQ-scaling in general does not follow as pointed in [12, 13]. The space momentum correlation of the parton system before hadronization requires detailed study of the dynamical evolution of the system [12, 14].

The level of NCQ-scaling also depends on the modelling of hadronization process. It is known that the finite width of the hadron internal wave function can lead to 10-20% violation of the v_2 NCQ-scaling [15]. In the extreme case where wave function is very wide, NCQ-scaling can be destroyed completely [16]. Since coalescence deals with $2, 3 \rightarrow 1$ processes, in general quarks or the coalescenced hadron need to be off mass shell [17], and the interactions with the surrounding medium are




FIG. 1: (Color online) Figure taken from [18] a) A compilation of v_2/n as function of p_T/n . b) The ratio to a common polynomial function.

necessary to neutralize their virtuality. Thus the constituent quarks should be treated as quasi-particles and there is no reason a priori to assume the momentum conservation in coalescence process (Eq.2). Our goal is to study the consequences of modifying this assumption.

If coalescence is a relevant hadronization mechanism, it should be valid at all momentum. However, from the data point of view, NCQ-scaling of the v_2 , when plotted in p_T , is not perfect [18], i.e. $v_2(p_T/n)/n$ is not a universal curve. This is illustrated by Fig.1, where the ratios of the scaled elliptic flow (v_2/n) for various hadrons to a combined polynomial fit are plotted as function of scaled transverse momentum (p_T/n). NCQ-scaling only works at 20% level at $p_T > 2$ GeV/c, then it breaks badly at $p_T < 2$ GeV/c. The breaking has a systematic dependence on the mass. It undershoots the v_2 values for light mesons such as pion and overshoots the v_2 for heavy baryons such as Ξ . Attempts have been made to reconcile the breaking of pion v_2 by including the resonance effect in the coalescence models [11, 19]. The agreement with data can be achieved qualitatively. We are not aware of any mechanisms within existing coalescence models that can accommodate the deviation of heavy baryons.

Recently it was pointed out by PHENIX Collaboration [20] that the constituent quark scaling of v_2 can be preserved at low p_T if hadron transverse kinetic energy $KE_T = m_T - m$ instead of p_T is used as a scaling variable. We duplicated the PHENIX finding in Fig.2. The deviation from the universal $v_2(KE_T/n)/n$ scaling curve is very small, less than 10% in all cases. Motivated by this observation, we modify the momentum conservation relation into kinetic energy conservation relation in the coalescence formula Eq.2:

$$W_M = \Phi_M(x_1 - x_2, p_1 - p_2) \delta(m_T - m_{T,1} - m_{T,2} - m + m_1 + m_2) \quad (6)$$




FIG. 2: (Color online) Figure taken from [20]: a) A compilation of v_2/n as function of KE_T/n . b) The ratio to a common polynomial function.

Without very detailed calculations, we can make the following remarks.

1. For the naive coalescence scenario, where only the co-moving q and $\bar{q}$ recombine into hadron, assuming spatially uniform quark density $n(x, p) = n(p)$ or flow anisotropy $v_{2,q}(x, p) = v_{2,q}(p)$, we obtain the usual NCQ-scaling relation for meson and baryon [2, 5], except it is in KE_T , instead of p_T .

$$v_{2,M}(KE_T) = \frac{2v_{2,q}}{1 + 2v_{2,q}^2} \bigg|_{\frac{KE_T}{2}} \approx 2v_{2,q}\left(\frac{KE_T}{2}\right) \quad (7)$$

$$v_{2,B}(KE_T) = \frac{3v_{2,q} + 3v_{2,q}^3}{1 + 6v_{2,q}^2} \bigg|_{\frac{KE_T}{3}} \approx 3v_{2,q}\left(\frac{KE_T}{3}\right) \quad (8)$$

We have confirmed these relations through a simple Monte-Carlo simulation of the naive coalescence process, where q or $\bar{q}$ are assumed to have mass of 350 MeV and an exponential spectra in m_T . In addition, similar to what is done in other recombination models, the transverse momentum of the combining quarks in x or y direction are required to be within 0.24 GeV/c of each other. In essence the coalescence formula Eq.1 becomes

$$\frac{d^3 N}{dp^3} = \int d^2 p_1 d^2 p_2 e^{-(m_{T,1} + m_{T,2})/A} \delta(KE_T - KE_{T,1} - KE_{T,2}) \Theta(\Delta_p - |p_{x,1} - p_{x,2}|) \Theta(\Delta_p - |p_{y,1} - p_{y,2}|) \quad (9)$$

where $A = 0.17$ GeV and $\Delta_p = 0.24$ GeV/c is the cut off of relative momentum between the two quarks. Since we focus on mid-rapidity, longitudinal momentum direction has been ignored in this study. Results of our calculations




FIG. 3: (Color online) a) The input quark $v_2(KE_T)$. b) The ratio of the scaled v_2 ($v_2(KE_T/n)/n$) for meson and baryon over quark v_2 (solid lines) and ratios of the scaled v_2 from Eq.7 and Eq.8 over quark v_2 (dashed lines).

are shown in Fig.3. The calculations are also compared with the analytical results given by Eq.7 and 8 in dashed lines. Clearly, the breaking of scaling due to the finite momentum spread in coalescence become significant only at $KE_T < 1$ GeV. Above 1 GeV, both curves approach the value indicated by Eq.7 and 8 (indicated by dashed lines). However, the split between meson and baryon in KE_T/n would not be discernable given the precision of the data shown by Fig.2. We noted that Eq.7 and Eq.8 are valid as long as the constituent quark spectra are exponential in m_T , independent of the quark mass.

2. The kinetic energy conservation condition is more than a simple improvement over the previous coalescence models only at low p_T . It represents a quantitatively different recombination condition at all p_T . At large p_T ($p_T \gg m$), coalescence condition Eq.6 becomes $p_T - p_{T,1} - p_{T,2} = m - m_1 - m_2$. It is different from momentum conservation, because in general $m \neq m_1 + m_2$. This can be clearly seen by comparing the bottom panels of Fig.1 and Fig.2. Going from p_T/n to KE_T/n , one effectively shift high p_T part of light meson v_2/n to the right and the low high p_T part of heavy baryon v_2/n to the left by an amount on the order of the hadron mass. However, the resulting scaled v_2 curves falls on top of each other.

3. In hydrodynamic model with Cooper-Fryer freeze

out [21], hadrons are statistically produced in the local co-moving frame with certain radial flow. At low p_T , the elliptic flow should be proportional to p_T^2 according to the a general analytical expansion [22]. The Buda-Lund hydro model [23] suggests a universal elliptic flow scaling : $v_2 = I_1(w)/I_0(w)$, $w \propto p_T^2/2m_T$, which, at low p_T (thus small w), reduce to $v_2 \approx 0.5w \propto p_T^2/2m = KE_T$. We also arrive this relation through a schematic derivation [24].

For any variable X and a linear relation between elliptic flow and X : $v_2 \propto X$, NCQ-scaling in X is a trivial additive artifact: $v_2/n \propto X/n$. It was realized by [13, 14] that either hydrodynamical mass ordering or NCQ-scaling of v_2 can results from quark coalescence. In our point of view, the coalescence condition Eq.6 provide a natural reconciliation between the statistical hadronization and hadronization via coalescence. Since v_2 is proportional to KE_T at low p_T , v_2 for various hadrons naturally fall on a universal curve in small KE_T independent of whether they are scaled by the number of constituent quark or not.

Due to the non-linear relation between p_T and KE_T , at low or intermediate p_T region where mass effect is important, NCQ-scaling can not hold in both p_T and KE_T simultaneously. At $p_T \gg m$, a schematic hydro calculation gives $v_2 \approx \tanh(\xi p_T)$ [25], where ξ depends on flow velocity. However, this relation gives neither NCQ-scaling in p_T nor NCQ-scaling in KE_T .

A complete model should be able to describe not only elliptic flow but also the particle ratio, single particle spectra, etc. Our study is rather limited in this regard. We do not attempt to describe any observables other than elliptic flow. Various hadronic effects after quark coalescence, such as the hadronic re-scattering and resonance decay, are ignored in current study. Our discussion is focused on the physical consequences when quark coalescence is extended to low p_T . We have demonstrated that results from a simple coalescence approach agree with the data fairly well in KE_T . The deviation of the pion v_2/n from p_T/n scaling is naturally resolved when plotted in KE_T/n variable, without the need to rely on the resonance effect. The fact that RHIC data indicate strong flavor dependence of the v_2 and strong violation of the NCQ-scaling of the v_2 in p_T suggests that KE_T scaling of the v_2 at intermediate p_T is a rather non-trivial observation. It suggests that the quark coalescence might be an important hadronization mechanism at low p_T ($p_T < 2$ GeV/c).

We thank M. Issah for pointing us to the Fig.2 in his talk [20]. Discussions with R. Fries, C. M. Ko are appreciated.

-
- [1] S. A. Voloshin, Nucl. Phys. A **715**, 379 (2003)
 [2] R. J. Fries, B. Muller, C. Nonaka and S. A. Bass, "Hadronization in heavy ion collisions: Recombination

- and fragmentation of Phys. Rev. Lett. **90**, 202303 (2003),
 Phys. Rev. C **68**, 044902 (2003)
 [3] V. Greco, C. M. Ko and P. Levai, Phys. Rev. Lett. **90**,

- 202302 (2003), Phys. Rev. C **68**, 034904 (2003)
- [4] R. C. Hwa and C. B. Yang, Phys. Rev. C **67**, 034902 (2003)
- [5] D. Molnar and S. A. Voloshin, Phys. Rev. Lett. **91**, 092301 (2003)
- [6] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 172301 (2003)
- [7] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 052302 (2004)
- [8] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 182301 (2003)
- [9] B. Muller, R. J. Fries and S. A. Bass, Phys. Lett. B **618**, 77 (2005)
- [10] C. B. Yang, J. Phys. G **32**, L11 (2006)
- [11] V. Greco and C. M. Ko, Phys. Rev. C **70**, 024901 (2004)
- [12] D. Molnar, arXiv:nucl-th/0512001.
- [13] S. Pratt and S. Pal, Nucl. Phys. A **749**, 268 (2005) [Phys. Rev. C **71**, 014905 (2005)]
- [14] D. Molnar, arXiv:nucl-th/0408044.
- [15] V. Greco and C. M. Ko, arXiv:nucl-th/0505061.
- [16] Z. w. Lin and C. M. Ko, Phys. Rev. Lett. **89**, 202302 (2002)
- [17] R. Scheibl and U. W. Heinz, Phys. Rev. C **59**, 1585 (1999)
- [18] M. Oldenburg [STAR Collaboration], arXiv:nucl-ex/0510026.
- [19] X. Dong, S. Esumi, P. Sorensen, N. Xu and Z. Xu, Phys. Lett. B **597**, 328 (2004)
- [20] A. Adare *et al.* [PHENIX Collaboration], arXiv:nucl-ex/0608033; M. Is-sah, talk in AGS User meetings, http://www.bnl.gov/rhic_ags/users_meeting/Workshops/5.asp, http://www.phenix.bnl.gov/phenix/WWW/publish/missah/RHIC-AGS-Users06/rhic-ags_missah.ppt
- [21] P. F. Kolb, J. Sollfrank and U. W. Heinz, Phys. Rev. C **62**, 054909 (2000)
- [22] P. Danielewicz, Phys. Rev. C **51**, 716 (1995)
- [23] M. Csanad, T. Csorgo and B. Lorstad, Nucl. Phys. A **742**, 80 (2004)
- [24] For a thermal source and at $p_T \ll mv_T$, where v_T is the radial flow velocity,
- $$f(p, x) \propto \exp\left(-\frac{\gamma_T(m_T - \vec{p}_T \cdot \vec{v}_T) \pm \mu}{T}\right) \quad (10)$$
- $$\propto \exp\left(-\frac{\gamma_T m_T}{T}\right) \propto \exp\left(-\frac{m \langle v(\phi) \rangle^2}{2T}\right)$$
- we have ignored the ϕ dependence of the T and μ . For this simple scenario, the elliptic flow would be,
- $$v_2 \approx \frac{m \left[\langle v \rangle_y^2 - \langle v \rangle_x^2 \right]}{8T} = \frac{p_T^2 \epsilon_v}{8mT} \quad (11)$$
- $\epsilon_v = \left(\langle v \rangle_y^2 - \langle v \rangle_x^2 \right) / \langle v \rangle^2$ is the velocity anisotropy and should to first order be proportional to spacial anisotropy [21]: $\epsilon_v = A\epsilon_x + O(p_T)$.
- [25] P. Huovinen, P. F. Kolb, U. W. Heinz, P. V. Ruuskanen and S. A. Voloshin, Phys. Lett. B **503**, 58 (2001)